

Supplementary Material

Green synthesis of 1,4-benzoxazin-3-one using a choline chloride based deep eutectic solvent

Ali Sharifi*, Marzieh Sharifi, M. Saeed Abaee, and Mojtaba Mirzaei

*Faculty of Organic Chemistry and Natural Products, Chemistry and Chemical Engineering
Research Center of Iran, Pajohesh Blvd., Tehran, Iran, 1496813151*

Email: sharifi@ccerci.ac.ir

Table of Contents

¹ H and ¹³ C NMR spectra of compounds.....	S2
--	----




Figure 1: ¹H NMR spectrum of benzoxazine 3fa

v




Figure 1: ¹³C NMR spectrum of benzoxazine 3fa




Figure 3: ¹H NMR spectrum of benzoxazine 3fc




Figure 4: ¹³C NMR spectrum of benzoxazine 3fc




Figure 5: ¹H NMR spectrum of benzoxazine 3ga




Figure 6: ¹³C NMR spectrum of benzoxazine 3ga




Figure7: ¹H NMR spectrum of benzoxazine 3gc




Figure8: ¹³C NMR spectrum of benzoxazine 3gc




Figure 9: ¹H NMR spectrum of benzoxazine 3hc




Figure 10: ¹³C NMR spectrum of benzoxazine 3hc