

Supplementary Material

An efficient and mild method for the synthesis of tetrazolo-fused-pyridine derivatives

Machireddy Rajesh Kumar Reddy^a, Arun Kumar^b, Deepak Kumar^b, and Ashok Kumar Yadav^{c*}

^a *Department of Energy Science and Engineering, DGIST, 333, Techno Jungang Daero, Hyeongpun-Eup,*

Dalseong-Gun, Daegu, 42988 South Korea


^b *School of Pharmaceutical Sciences, Shoolini University of Biotechnology and Management Sciences, India*

^c *University Institute of Pharmaceutical Sciences, Panjab University, India*


Email: ashoky@pu.ac.in

Table of Contents


| | |
|--------------------------------------------------------------------------------------------------|----|
| ¹ H and ¹³ C NMR Spectra of Tetrazolo-fused Pyridines (2a-g)..... | S2 |
|--------------------------------------------------------------------------------------------------|----|




¹H NMR spectra of tetrazolo[1,5-a]pyridine (2a)




¹³C NMR spectra of tetrazolo[1,5-a]pyridine (2a)




¹H NMR spectra of 7-methoxytetrazolo[1,5-*a*]pyridine (2b)




¹³C NMR spectra of 7-methoxytetrazolo[1,5-*a*]pyridine (2b)




¹H NMR spectra of 7-Methyltetrazolo[1,5-*a*]pyridine (2c)




¹³C NMR spectra of 7-Methyltetrazolo[1,5-*a*]pyridine (2c)




¹H NMR spectra of Tetrazolo[1,5-*a*]pyridine-7-carbonitrile (2d)




¹³C NMR spectra of Tetrazolo[1,5-*a*]pyridine-7-carbonitrile (2d)




¹H NMR spectra of Tetrazolo[1,5-a]quinoline (2e)




¹³C NMR spectra of Tetrazolo[1,5-a]quinoline (2e)




¹H NMR spectra of 7-Methoxytetrazolo[1,5-*a*]quinoline (2f)



¹³C NMR spectra of 7-Methoxytetrazolo[1,5-*a*]quinoline (2f)



¹H NMR spectra of Tetrazolo[5,1-a]isoquinoline (2g)



¹³C NMR spectra of Tetrazolo[5,1-a]isoquinoline (2g)